arXiv:hep-ph/0611045v1 3 Nov 2006

BARI-TH/06-552

The three flavor LOFF phase of QCD

Nicola D. Ippolito*
Università di Bari, I-70126 Bari, Italy and
I.N.F.N., Sezione di Bari, I-70126 Bari, Italy

(Dated: February 2, 2008)

We explore the Larkin-Ovchinnikov-Fulde-Ferrell (LOFF) phase of QCD with three flavors, using a Ginzburg-Landau expansion of the free energy, and a NJL point-like four-fermion interaction, with the quantum numbers of single-gluon exchange.

PACS numbers: 12.38.Aw, 12.38.Lg

At high quark densities and small temperatures quarks are expected to form Cooper pairs, because of the attractive interaction in the color antisymmetric channel. This gives rise to Color Superconductivity, see [1, 2] and for reviews [3, 4]. At asymptotically high densities with three flavors the Color Flavor Locking (CFL) phase [5], characterized by a spinless color- and flavor-antisymmetric diquark condensate, is the energetically favored phase. When the density decreases to values probably achieved in the core of compact stars ($\mu \sim 400 - 500$ MeV) one cannot neglect the strange quark mass and the differences in the quark chemical potentials $\delta\mu$. Various phases have been proposed for this region of the QCD phase diagram. Among these, the gapless gCFL phase [6] was considered for some time as the most suitable candidate. It was however realized that imaginary gluon Meissner masses, see [7], produce instabilities in this phase, and in general in the gapless phases, see [8].

The Larkin-Ovchinnikov-Fulde-Ferrell (LOFF) [9] phase is an inhomogeneous superconductive phase in which quark pairs have non-zero total momentum; it turns out to be favored, for appropriate values of $\delta\mu$, over the homogeneous superconducting phase and the normal non-superconducting one. The possibility of a LOFF phase in QCD with two flavors has been extensively discussed in the last half-decade, see [10] and for a review [11].

The three flavor LOFF phase has been first considered in [12], using a Ginzburg-Landau (GL) expansion of the free energy, and subsequently it was found that such a phase is chromomagnetically stable [13]. The validity of the G-L approximation for this phase has recently been tested [14], and a study of the corrections due to the finite chemical potentials has been completed in [15]. Recently a work has been performed [16] in which the authors consider the possibility of structures formed by more than one plane wave.

Formalism and results

We start with the Lagrangean density for three flavor ungapped quarks

$$\mathcal{L} = \bar{\psi}_{i\alpha} \left(i \, \not{D}_{ij}^{\alpha\beta} - M_{ij}^{\alpha\beta} + \mu_{ij}^{\alpha\beta} \gamma_0 \right) \psi_{\beta j} \ . \tag{1}$$

$M_{ij}^{\alpha\beta} = \delta^{\alpha\beta} \, \mathrm{diag}(0, 0, M_s)$ is the mass matrix and $D_{ij}^{\alpha\beta} = \partial \delta^{\alpha\beta} \delta_{ij} + ig A_a T_a^{\alpha\beta} \delta_{ij}$; $\mu_{\alpha\beta}^{ij}$ is a diagonal color-flavor matrix depending in general on μ (the average quark chemical potential), μ_e (the electron chemical potential), and μ_3, μ_8, related to color [6]. We do not require color neutrality and we work in the approximation $\mu_3 = \mu_8 = 0$, which is justified by the results of Ref.[15], so we write $\mu_{ij}^{\alpha\beta} = (\mu \delta_{ij} - \mu_e Q_{ij}) \delta^{\alpha\beta} = \mu_i \, \delta_{ij} \delta^{\alpha\beta}$.
Treating the strange quark mass at the leading order in the $1/\mu$ expansion we have

$$\mu_u = \mu - \frac{2}{3}\mu_e \ , \ \mu_d = \mu + \frac{1}{3}\mu_e \ , \ \mu_s = \mu + \frac{1}{3}\mu_e - \frac{M_s^2}{2\mu} \ . \tag{2}$$

Moreover we make use of the High Density Effective Theory (HDET), see [17] and, for a review, [4].

The fundamental pairing ansatz for the condensate is

$$< \psi_{i\alpha} \, C \, \gamma_5 \, \psi_{\beta j} >= \sum_{I=1}^{3} \Delta_I(\mathbf{r}) \, \epsilon^{\alpha\beta I} \, \epsilon_{ijI} \tag{3}$$

---

*Electronic address: nicola.ippolito@ba.infn.it with

$$\Delta_I(\mathbf{r}) = \Delta_I \exp\left(2i\mathbf{q_I}\cdot\mathbf{r}\right)\ , \tag{4}$$

where $2\mathbf{q_I}$ is the pair momentum. The momenta $\mathbf{q_1}$, $\mathbf{q_2}$, $\mathbf{q_3}$ and the gap parameters $\Delta_1(\mathbf{r})$, $\Delta_2(\mathbf{r})$, $\Delta_3(\mathbf{r})$ correspond respectively to $d-s$, $u-s$ and $u-d$ pairing.

As usual, the norms $|\mathbf{q_I}|$ are derived minimizing the free energy, while the directions have to be determined by a chrystallographic analysis. In [12] only structures with the $\mathbf{q_I}$ on the same axis have been studied, while in [16] more complicate structures are analyzed.

The GL approximation is implemented by an expansion of the anomalous propagator S_{21}

$$S_{21} = S_0^{22}\Delta^* S_0^{11} + S_0^{22}\Delta^* S_0^{11}\Delta S_0^{22}\Delta^* S_0^{11} + O(\Delta^5) \tag{5}$$

involved in the gap equation, as obtained by the Schwinger-Dyson equation in the HDET formalism:

$$\Delta^*_{AB}(\mathbf{r}) = i\,3G\,V^\mu\tilde{V}^\nu \sum_{C,D=1}^{9} h^*_{AaC}h_{DbB}\int\frac{d\,\mathbf{n}}{4\pi}\int\frac{d^3\,\ell}{(2\pi)^3}\int\frac{dE}{2\pi}\,S_{21}(E,\ell)_{CD}\,g_{\mu\nu}\,\delta_{ab}\ . \tag{6}$$

In this way the free energy GL expansion assumes the form

$$\Omega = \Omega_n + \sum_{I=1}^{3}\left(\frac{\alpha_I}{2}\,\Delta_I^2 \ + \ \frac{\beta_I}{4}\,\Delta_I^4 \ + \ \sum_{J\neq I}\frac{\beta_{IJ}}{4}\,\Delta_I^2\Delta_J^2\right) \ + \ O(\Delta^6) \tag{7}$$

with

$$\Omega_n = -\frac{3}{12\pi^2}\left(\mu_u^4+\mu_d^4+\mu_s^4\right) - \frac{\mu_e^4}{12\pi^2}\ . \tag{8}$$

The coefficients α_I, β_I and β_{IJ} can be found in [12]. Electric neutrality is obtained by imposing the condition

$$-\frac{\partial\Omega}{\partial\mu_e} = 0\ . \tag{9}$$

.

In [12] the energetically favored solution has $\Delta_1 = 0$, $\Delta_2 = \Delta_3$ and $\mathbf{q_2}$, $\mathbf{q_3}$ parallel. Figure 1 shows the competition among different color superconducting phases, comprising the three flavor LOFF phase considered in [12].



FIG. 1: Free energy differences $\Omega_{LOFF} - \Omega_{norm}$ in units of 10^6 MeV^4 plotted versus M_s^2/μ (in MeV) for various QCD phases.

In Ref.[16] another plot is presented, showing the free energies of more complicate crystal structures. In particular, there are two structures with free energy smaller than the simple ansatz of [12], but the gaps involved in this phases are rather large, so the GL approximation must be properly tested.

In conclusion, we have shown that the three flavor LOFF phase, due to its chromomagnetic stability, is a serious candidate for the true vacuum at moderate densities, and therefore it could be present in the core of compact stars (see [18] for a preliminary study).

Acknowledgments

I wish to thank G. Nardulli and M. Ruggieri, for their fundamental importance upstream this work; D. Blaschke and BLTP-JINR for everything concerning the DM 2006 school; R. Anglani and all the other participants to DM 2006 for two really useful and funny weeks.

---

[1] J. C. Collins and M. J. Perry, Phys. Rev. Lett. **34** (1975) 1353; B. Barrois, Nucl. Phys. **B129** (1977) 390; S. Frautschi, Proceedings of workshop on hadronic matter at extreme density, Erice 1978; D. Bailin and A. Love, Phys. Rept. **107** (1984) 325.

[2] M. G. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. B **422**, 247 (1998) [arXiv:hep-ph/9711395]; R. Rapp, T. Schäfer, E. V. Shuryak and M. Velkovsky, Phys. Rev. Lett. **81**, 53 (1998) [arXiv:hep-ph/9711396]; D. T. Son, Phys. Rev. D **59**, 094019 (1999) [arXiv:hep-ph/9812287]; R. D. Pisarski and D. H. Rischke, Phys. Rev. D **61**, 074017 (2000) [arXiv:nucl-th/9910056].

[3] K. Rajagopal and F.Wilczek,in Handbook of QCD, M. Shifman ed. (World Scientific 2001), [arXiv:hep-ph/0011333]; M. G. Alford, Ann. Rev. Nucl. Part. Sci. **51** (2001) 131 [arXiv:hep-ph/0102047]. T. Schafer, arXiv:hep-ph/0304281.

[4] G.Nardulli, Riv. Nuovo Cim. **25N3** (2002) 1 [arXiv:hep-ph/0202037].

[5] M. G. Alford, K. Rajagopal and F. Wilczek, Nucl. Phys. B **537**, 443 (1999) [arXiv:hep-ph/9804403].

[6] M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev. Lett. **92**, 222001 (2004) [arXiv:hep-ph/0311286]; M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev. D **71**, 054009 (2005) [arXiv:hep-ph/0406137]. K. Fukushima, C. Kouvaris and K. Rajagopal, Phys. Rev. D **71**, 034002 (2005) [arXiv:hep-ph/0408322].

[7] R. Casalbuoni, R. Gatto, M. Mannarelli, G. Nardulli and M. Ruggieri, Phys. Lett. B **605**, 362 (2005) [Erratum-ibid. B **615**, 297 (2005)] [arXiv:hep-ph/0410401]; K. Fukushima, Phys. Rev. D **72**, 074002 (2005) [arXiv:hep-ph/0506080]; M. Alford and Q. h. Wang, J. Phys. G **31**, 719 (2005) [arXiv:hep-ph/0501078]; K. Fukushima, arXiv:hep-ph/0510299.

[8] M. Huang and I. A. Shovkovy, Phys. Rev. D **70**, 051501 (2004) [arXiv:hep-ph/0407049]. M. Huang and I. A. Shovkovy, Phys. Rev. D **70**, 094030 (2004) [arXiv:hep-ph/0408268].

[9] A. I. Larkin and Yu. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47** (1136) 1964 (Sov. Phys. JETP **20** (1965) 762); P.Fulde and R. A. Ferrell, Phys. Rev. **135** (1964) A550.

[10] M. G. Alford, J. A. Bowers and K. Rajagopal, Phys. Rev. D **63**, 074016 (2001) [arXiv:hep-ph/0008208]; J. A. Bowers, J. Kundu, K. Rajagopal and E. Shuster, Phys. Rev. D **64**, 014024 (2001) [arXiv:hep-ph/0101067]; J. A. Bowers and K. Rajagopal, Phys. Rev. D **66**, 065002 (2002) [arXiv:hep-ph/0204079]; R. Casalbuoni, M. Ciminale, M. Mannarelli, G. Nardulli, M. Ruggieri and R. Gatto, Phys. Rev. D **70**, 054004 (2004) [arXiv:hep-ph/0404090].

[11] R. Casalbuoni and G. Nardulli, Rev. Mod. Phys. **76**, 263 (2004) [arXiv:hep-ph/0305069].

[12] R. Casalbuoni, R. Gatto, N. Ippolito, G. Nardulli and M. Ruggieri, Phys. Lett. B **627**, 89 (2005) [Erratum-ibid. B **634**, 565 (2006)] [arXiv:hep-ph/0507247].

[13] M. Ciminale, G. Nardulli, M. Ruggieri and R. Gatto, Phys. Lett. B **636**, 317 (2006) [arXiv:hep-ph/0602180].

[14] M. Mannarelli, K. Rajagopal and R. Sharma, arXiv:hep-ph/0603076.

[15] R. Casalbuoni, M. Ciminale, R. Gatto, G. Nardulli and M. Ruggieri, arXiv:hep-ph/0606242.

[16] K. Rajagopal and R. Sharma, arXiv:hep-ph/0605316, K. Rajagopal and R. Sharma, arXiv:hep-ph/0606066.

[17] D. K. Hong, Phys. Lett. B **473**, 118 (2000) [arXiv:hep-ph/9812510]. D. K. Hong, Nucl. Phys. B **582**, 451 (2000) [arXiv:hep-ph/9905523]; S. R. Beane, P. F. Bedaque and M. J. Savage, Phys. Lett. B **483**, 131 (2000) [arXiv:hep-ph/0002209]; R. Casalbuoni, R. Gatto, G. Nardulli and M. Ruggieri, Phys. Rev. D **68**, 034024 (2003) [arXiv:hep-ph/0302077].

[18] R. Anglani, G. Nardulli, M. Ruggieri, M. Mannarelli, Phys. Rev. D **74** 074005 [arXiv:hep-ph/0607341]